

# Nucleophilic trifluoromethylation of some polycyclic ketones <sup>#</sup>

Jarosław Romański\* and Grzegorz Młostoń

*University of Łódź, Department of Organic and Applied Chemistry,  
Section of Heteroorganic Compounds, Narutowicza 68, 90-136 Łódź, Poland  
E-mail: [romanski@uni.lodz.pl](mailto:romanski@uni.lodz.pl)*

**Dedicated to Prof. Jan Epszajn on the occasion of his 75<sup>th</sup> birthday**

---

## Abstract

Ruppert's reagent ( $\text{CF}_3\text{-SiMe}_3$ ) was used in the reaction with derivatives of pentacyclo-[5.4.0.0<sup>2,6</sup>.0<sup>3,10</sup>.0<sup>5,9</sup>]undecane-8,11-dione ('cage' dione) in the presence of dry CsF to yield trifluoromethyl *O*-silylated products. Subsequent acidic hydrolysis gave the corresponding hydroxy derivatives. In the case of the 'cage' dione the transannular cyclization leading to oxahexacyclic (*O*-bridged) product was observed.

**Keywords:** Trifluoromethylation, polycyclic ketones, nucleophilic addition, transannular cyclization

---

## Introduction

The chemistry of fluorinated compounds has been extensively explored in recent decades. It is well established that introduction of the trifluoromethyl group into a known substance results in significant changes in the chemical, physical and biological properties.<sup>1</sup> Direct trifluoromethylation of aromatic compounds is mostly based on the reactions of *in situ* generated  $\cdot\text{CF}_3$  radical.<sup>2</sup> The non-direct trifluoromethylation with  $\text{SF}_4$  used as a source of fluorine atoms has been developed by Dmowski.<sup>3</sup> On the other hand, there are known methods that explore an ionic  $\text{CF}_3$ -moiety, *e.g.* the sulfonium salt of type **1** is an efficient source of the trifluoromethyl cation (Scheme 1).<sup>4</sup> In recent times, nucleophilic trifluoromethylation has been widely explored for introduction of the  $\text{CF}_3$ -group using carbonyl substrates such as aldehydes, ketones or esters<sup>5</sup> and trimethyl(trifluoromethyl)silane (**3**) known as the Ruppert's reagent.<sup>6</sup>



---

<sup>#</sup> For the first time presented at 14<sup>th</sup> European Symposium on Fluorine Chemistry, Poznań, Poland, July 11-16, 2004, Book of Abstracts, A-P-75, 164.



R = e.g.: Br, I, SiEt<sub>3</sub>, ZnX, Cu, COOMe


X = H or Hal



**Scheme 1**

In the original protocol, tetrabutylammonium fluoride (TBAF) was used as a catalyst in the reaction of **3** with carbonyl compounds yielding the corresponding trifluoromethylated alcohols via *O*-silylated derivatives.<sup>7</sup> At present, cesium fluoride seems to be a catalyst of choice.<sup>5a</sup> However recently, an efficient application of the new catalysts in nucleophilic trifluoromethylation was described.<sup>8</sup>

The relatively easily available polycyclic ‘cage’ ketones **6** and **9** can be used as building blocks for the preparation of more complex systems, including hosts in supramolecular chemistry.<sup>9</sup> Adamantanone (**2**), which belongs to the polycyclic ‘cage’ structures, reacts easily with **3** yielding trimethylsilyl derivative **4** which after acidic hydrolysis can be easily converted to 2-(trifluoromethyl)adamantan-2-ol (**5**) (Scheme 2).<sup>10</sup>




**Scheme 2**

In the case of the dione **9** reactions with primary amines, are reported to yield the corresponding monoimine derivatives which upon treatment with reducing agents (NaBH<sub>4</sub>, LiAlH<sub>4</sub>) undergo a transannular cyclization via formation of the O-bridge.<sup>11</sup> Reactivities of **6** and **9** have been studied in our group for some years and their ability to undergo transannular cyclization was observed in reactions with diverse O- and S-nucleophiles.<sup>12</sup> The aim of the

present study was to examine their reactions with Ruppert's reagent **3** and to elaborate a straightforward protocol for preparation of trifluoromethylated polycyclic alcohols.

## Results and Discussion

The experiment with **3** and **6** was performed in dry 1,2-dimethoxyethane (DME) in the presence of catalytic amount of cesium fluoride (dried at 200-300 °C). The expected *O*-silylated product **7** was smoothly obtained and subsequent hydrolysis followed by crystallization from hexane led to a colorless solid identified as alcohol **8** (Scheme 3).




**Scheme 3**


The reaction occurs stereoselectively and the trifluoromethyl anion approaches the carbonyl group from the *exo*-face exclusively. As a result, the final product **8** contained the hydroxy group placed at the *endo* position as evidenced by the X-ray study.<sup>13</sup>

The reaction of **3** with **9** was carried out under analogous conditions. Depending on the molar ratio of substrates dione **9** was expected to react either with one or two equivalents of CF<sub>3</sub>-SiMe<sub>3</sub> yielding mono- or bis-trifluoromethylated alcohols **10** or **11**, respectively. The first attempted reaction of **9** with an equimolar amount of **3** was expected to yield the hydroxyketone **10** (Scheme 4).

After addition of CF<sub>3</sub>-SiMe<sub>3</sub> an exothermic reaction started and after 1.5 h the conversion was complete. The initially formed trimethylsilyl derivative was hydrolyzed without isolation and crude product was purified by crystallization from hexane to give colorless crystals with a narrow melting point. The second experiment with two equivalents of CF<sub>3</sub>-SiMe<sub>3</sub> should have led to **11**, but unexpectedly the isolated product was identical with the substance obtained in the first entry; this result was confirmed by the absence of mixed melting point depression and by comparison of spectroscopic data. The IR-spectrum did not show the absorption band of the C=O group suggesting that a transannular process took place. In the <sup>13</sup>C-NMR spectra the characteristic signals appeared at 89.3 (q, <sup>2</sup>J<sub>C-F</sub> = 34 Hz) and 120.0 (s, O-C-O) ppm, respectively. The MS-Cl exhibited a molecular peak at m/z 262 [M+1+NH<sub>3</sub>]<sup>+</sup>. Finally, the structure of trifluoromethylated hemiacetal **13** was unambiguously confirmed by the X-ray single crystal diffraction analysis (Fig.1).<sup>13</sup>




### Scheme 4



**Figure 1.** ORTEP-plots for compounds 13 and 21.<sup>13</sup>

Noteworthy, the transannular cyclization under trifluoromethylation conditions was previously reported for the reaction of a 1,5-dione (*i.e.* bicyclo[3.3.1]nonane-2,6-dione) but in this case the successful formation of the product of bis-trifluoromethylation was reported.<sup>14</sup>

According to an other report the reaction of 1,4-dione (*i.e.* hexane-2,5-dione) with one or two equivalents of **3** did not afford the product of transannular cyclization.<sup>15</sup> Recently, we have found that **9** easily reacts at low temperatures with amines as a nucleophilic agent and the formation of transannular cyclization products was established,<sup>16</sup> instead of already reported aminoalcohol derivatives.<sup>11</sup>




**Scheme 5**

The formation of **13** strongly supports the mechanistic pathway of the CF<sub>3</sub>-group addition to the carbonyl group formulated by Prakash *et al.*<sup>17</sup> The first step of the reaction is the replacement of the CF<sub>3</sub>-group by the fluorine anion and trifluoromethide anion (F<sub>3</sub>C<sup>-</sup>) attacks the carbonyl group to form the salt of type **14**. Apparently, the following transannular cyclization leading to **15** is faster than the addition of another molecule of **3** leading to the intermediate **17** which could give hydroxyketone **10**. In the next step, reaction with another molecule of **3** generates a transient species **16** bearing pentacoordinated silicon atom. This intermediate rapidly transfers F<sub>3</sub>C-group to the next molecule of **9** and the formation of the *O*-silylated acetal **12** is observed (Scheme 5).


In an extension of the above presented study, the C-C *back*-bond in the dione **9** was selectively reduced with zinc dust in refluxing glacial acetic acid yielding the ‘released’ ketone **18** (tetracyclo[6.3.0.0<sup>4,11</sup>.0<sup>5,9</sup>]undecane-2,7-dione).<sup>18</sup> This ketone was treated with **3** in order to

examine the limitation of transannular cyclization in this less strained structure. The distance between the two carbonyl groups, as evidenced by the X-ray study, is larger in comparison with ‘cage’ dione **9**.<sup>19,21b</sup> The product isolated after reaction with **3** and subsequent hydrolysis exhibited no C=O absorption, indicating the formation of hemiacetal **19**; additional spectroscopic data are comparable with those registered for compound **12**.



**Scheme 6**

In another experiment the imine **20** prepared in the reaction of aniline with **9**,<sup>20</sup> was treated with **3** and in this case the formation of silylated ether **21** was observed. The transannular process has not been observed and even with an excess of **3** the C=N-group did not react. The X-ray analysis<sup>13</sup> showed that the  $\text{CF}_3$ -group is placed at *exo*-position (Fig.1).



**Scheme 7**

In summary, the synthesis of new derivatives of trifluoromethylated polycyclic compounds **8**, **13** and **19** using Ruppert’s reagent was presented. Transannular cyclization led to hemiacetals **13** and **19**, but it was not observed in the reaction with monoimine **20**. In all cases the trifluoromethylation process proceeded stereoselectively and the trifluoromethyl group is placed at the *exo* position, exclusively.

## Experimental Section

**General Procedures.** Melting points were determined in a capillary by using a MelTemp 2 apparatus and are uncorrected. IR spectra were obtained using a NEXUS FT-IR apparatus. MS spectra were obtained using a Varian MAT-112S spectrometer, which was operated in the CI mode.  $^1\text{H}$ -,  $^{13}\text{C}$ - and  $^{19}\text{F}$ -NMR spectra were recorded with a Bruker 300 MHz in  $\text{CDCl}_3$  using

TMS ( $\delta = 0$  ppm) or  $\text{CFCl}_3$ , respectively, as an internal standards. Elemental analysis and HRMS (Finnigan MAT95) data were obtained at Polish Academy of Sciences.

### Starting materials

Trimethyl(trifluoromethylhyl)silane (**3**), pentacyclo[5.4.0.0<sup>2,6</sup>.0<sup>3,10</sup>.0<sup>5,9</sup>]undecane-8,11-dione (**9**), cesium fluoride and dry dimethoxyethane (DME) are commercialy avialable. Pentacyclo[5.4.0.0<sup>2,6</sup>.0<sup>3,10</sup>.0<sup>5,9</sup>]undecane-8-one<sup>21</sup> (**6**) and tetracyclo[6.3.0.0<sup>4,11</sup>.0<sup>5,9</sup>]undecane-2,7-dione<sup>18</sup> (**18**) were obtained according to described procedures.

### Reaction of Ketones **6**, **9** and **18** with Ruppert's Reagent - General Procedure

The corresponding ketone **6**, **9** or **18** (1 mmol) was dissolved in dry DME (3 ml) and to the solution catalytic amount of CsF (approx. 5 mg, dried at 200-300 °C) was added under argon atmosphere. The mixture was protected against the moisture. Trimethyl(trifluoromethylhyl)silane (**3**) (1.1 mmol) was added in small portions through the septum and the mixture was stirred for 1.5 h at ambient temp. After this time 4N HCl (1 ml) was added and stirring was continued for an additional 1 h. Then the reaction mixture was diluted with water (5 ml) and extracted with dichloromethane (3x5 ml). Combined organic layers were dried over anhydrous magnesium sulfate. The solvent was removed in *vacuo* and crude products were purified by crystallization.

**8-(Trifluoromethyl)pentacyclo[5.4.0.0<sup>2,6</sup>.0<sup>3,10</sup>.0<sup>5,9</sup>]undecane-8-ol (**8**).** mp 126–127 °C (hexane), yield 60 % (0.14 g, 0.6 mmol). <sup>1</sup>H NMR ( $\text{CDCl}_3$ )  $\delta$ : 1.06-1.13(m, 1H), 1.24(AB,  $J_{\text{AB}}=11$  Hz, 1H), 1.70(AB,  $J_{\text{AB}}=11$  Hz, 1H), 1.97(br. s, 1H, OH), 2.30-2.39(m, 2H), 2.40-2.68(m, 6H), 2.83-2.86(s, 1H). <sup>13</sup>C NMR ( $\text{CDCl}_3$ )  $\delta$ : 29.2(t), 34.7(t), 36.0(d), 38.3(d), 41.7(d), 42.1(d), 43.2(d), 43.5(d), 45.7(d), 46.3(d), 82.3(q,  $^2J_{\text{C-F}}=27$  Hz), 126.4(q,  $^1J_{\text{C-F}}=285$  Hz). <sup>19</sup>F NMR ( $\text{CDCl}_3$ )  $\delta$ : -76.4(s). IR (KBr)  $\text{cm}^{-1}$ : 3346vs, 2978s, 2867s, 1454w, 1393m, 1308s, 1288s, 1264s, 1158vs, 1143vs, 1127s, 1080s, 1026s, 967m, 700s. Anal. Calcd for  $\text{C}_{12}\text{H}_{13}\text{F}_3\text{O}$ : C 62.60; H 5.69. Found: C 62.61; H 5.57. MS (m/z): 248 ( $\text{M}+1+\text{NH}_3$ )<sup>+</sup>

**5-(Trifluoromethyl)-4-oxahexacyclo[5.4.1.0<sup>2,6</sup>.0<sup>3,10</sup>.0<sup>5,9</sup>.0<sup>8,11</sup>]dodecane-3-ol (**13**).** mp 116–117 °C (methanol), yield 65 % (0.16 g, 0.65 mmol). <sup>1</sup>H NMR ( $\text{CDCl}_3$ )  $\delta$ : 1.59(AB,  $J_{\text{AB}}=11$  Hz, 1H), 1.95(AB,  $J_{\text{AB}}=11$  Hz, 1H), 2.61-2.81(m, 6H), 2.83-3.11(m, 2H). <sup>13</sup>C NMR ( $\text{CDCl}_3$ )  $\delta$ : 41.9(d), 42.0(d), 43.4(d), 43.5(t), 45.3(d), 45.5(d), 47.1(d), 55.3(d), 57.0(d), 89.3(q,  $^2J_{\text{C-F}}=34$  Hz), 120.0(s), 124.7(q,  $^1J_{\text{C-F}}=275$  Hz). <sup>19</sup>F NMR ( $\text{CDCl}_3$ )  $\delta$ : -80.0(s). IR (KBr)  $\text{cm}^{-1}$ : 3348vs, 2993vs, 2976vs, 2881m, 1722m, 1396s, 1351s, 1307m, 1191vs, 1176vs, 1138s, 1035s, 861m, 715m. Anal. Calcd for  $\text{C}_{12}\text{H}_{11}\text{F}_3\text{O}_2$ : C 59.02; H 4.54. Found: C 59.29; H 4.21. MS (m/z): 262 ( $\text{M}+1+\text{NH}_3$ )<sup>+</sup>.

**7-(Trifluoromethyl)-12-oxapentacyclo[6.3.1.0<sup>2,6</sup>.0<sup>3,10</sup>.0<sup>5,9</sup>]dodecane-1-ol (**19**).** mp 90-92 °C (hexane), yield 57 % (0.14 g, 0.57 mmol). <sup>1</sup>H NMR ( $\text{CDCl}_3$ )  $\delta$ : 1.65-1.79(m, 2H), 1.92-2.12(m, 4H), 2.23-2.61(m, 5H), 2.78-3.15(m, 2H). <sup>13</sup>C NMR ( $\text{CDCl}_3$ )  $\delta$ : 35.5(t), 37.7(t), 41.7(d), 42.6(d), 43.3(t), 47.8(d), 49.3(d), 55.2(d), 58.9(d), 89.5(q,  $^2J_{\text{C-F}}=31$  Hz), 117.3(s), 125.8(q,  $^1J_{\text{C-F}}=278$  Hz). IR (KBr)  $\text{cm}^{-1}$ : 3399brs, 2971s, 2947m, 1377m, 1388s, 1299s, 1219s, 1170vs, 1145vs, 1119s,

1102s, 1062s, 1042s, 1012m, 947m. HRMS: Calcd. for C<sub>12</sub>H<sub>13</sub>F<sub>3</sub>O<sub>2</sub>: 246.0868; Found: 246.0863. MS-EI (m/z): 246 (M<sup>+</sup>, 32), 224(100), 186(72), 91(50).

**Phenyl(11-trifluoromethyl-11-trimethylsilyloxypentacyclo[5.4.0.0<sup>2,6</sup>.0<sup>3,10</sup>.0<sup>5,9</sup>]undec-8-ylidene)amine (21).** Dione **9** (250 mg, 1.15 mmol) was dissolved in THF (10 ml) and cooled down in ice-bath. To this solution aniline (0.5 ml, excess) was added dropwise and after approx. 30 min. white precipitate was filtered off. This half-product was suspended in benzene and dehydrated in boiling toluene using Dean-Stark apparatus. After 1h toluene was removed in *vacuo* and obtained imine **20** (200 mg, 0.80 mmol, 70%) was dissolved in dry DME (3 ml). To this solution catalytic amount of CsF (approx. 5 mg) was added under argon atmosphere and the mixture was protected against the moisture. The Ruppert's reagent **3** (1.0 mmol) was added in small portions through the septum and the mixture was stirred for 1.5 h at ambient temp. After this time water (1 ml) was added and stirring was continued for additional 1 h. Then the reaction mixture was diluted with water (5 ml) and extracted with dichloromethane (3x5 ml). Combined organic layers were dried over anhydrous magnesium sulfate. The solvent was removed in *vacuo* and crude product was purified by chromatography on preparative plates (SiO<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub>/hexane, 1:4). Obtained product **21** was recrystallized from MeOH to give colorless crystals (150 mg, 0.38 mmol, 48%), mp 140-142 °C.

<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 0.14(s, 9H), 1.50(AB, J<sub>AB</sub>=11Hz, 1H), 1.89(AB, J<sub>AB</sub>=11Hz, 1H), 2.46-2.96(m, 8H), 6.92-7.11(m, 3H), 7.26-7.34(m, 2H). <sup>13</sup>C NMR (CDCl<sub>3</sub>) δ: 33.3(d), 37.4(t), 38.9(d), 41.4(d), 42.2(t), 50.7(d), 51.1(d), 82.7(q, <sup>2</sup>J<sub>C-F</sub> = 27 Hz), 122.4(d), 124.0(d), 126.9(q, <sup>1</sup>J<sub>C-F</sub>=286 Hz), 128.3(d), 149.7(s), 178.8(s). <sup>19</sup>F NMR (CDCl<sub>3</sub>) δ: -75.2(s). IR (KBr) cm<sup>-1</sup>: 2997s, 2898m, 2870m, 1672m 1604m, 1596m, 1488m, 1406m, 1339s, 1307m, 1254vs, 1159vs, 1140vs, 1042s, 886s, 864s, 846s, 756m 701m. Anal. Calcd for C<sub>21</sub>H<sub>24</sub>F<sub>3</sub>NOSi: C 64.43; H 6.18. Found: C 64.18; H 6.07. MS (m/z): 392 (M+1)<sup>+</sup>.

## Acknowledgements

Authors thank the Rector of University of Lodz for financial support (Grant nr 505/712).

## References

1. (a) Olah, G. A.; Chambers, R. P.; Prakash, G. K. S. *Synthetic Fluorine Chemistry*; J. Wiley & Sons: New York, 1992. (b) Smart, B. E. *Organofluorine Chemistry, Principles and Commercial Applications*; Plenum Press: New York, 1994, pp 57-88. (c) *Methoden der organischen Chemie* (Houben-Weyl); Vol. E10a-b; Baasner, B.; Hagemann, H.; Tatlow, J. C.; Eds.; Thieme-Stuttgart: 1999/2000, Vol. E10a-b.
2. McClinton, M. A.; McClinton, D. A. *Tetrahedron* **1992**, *48*, 6555.

3. Dmowski, W. In *Methoden der organischen Chemie* (Houben-Weyl); Baasner, B.; Hagemann, H, Eds.; Thieme-Stuttgart: 1999, Vol. E10a, pp 321-395.
4. Umemoto, T.; Ishihara, S. *Tetrahedron Lett.* **1990**, *31*, 3579.
5. (a) Singh, R. P.; Shreeve, J. M. *Tetrahedron*, **2000**, *56*, 7613. (b) Prakash, G. K. S.; Mandal M. *J. Fluorine Chem.* **2001**, *112*, 123. (c) Kim, J.; Shreeve, J. M. *Org. Biomol. Chem.* **2004**, *2*, 2728. (d) Song, J. J.; Tan, Z.; Reeves, J. T.; Gallou, F.; Nathan, K.; Yee, N. K.; Senanayake, C. H. *Org. Lett.*, **2005**, *7*, 2193.
6. (a) Ruppert, I.; Schlich, K.; Volbach, W. *Tetrahedron Lett.* **1984**, *24*, 2195. (b) Thayer, A. M. *Chem. Eng. News*, **2006**, *84*, 15.
7. Prakash, G. K. S.; Krishnamurti, R.; Olah, G. A. *J. Am. Chem. Soc.* **1989**, *111*, 393.
8. Prakash, G. K. S.; Panja, C.; Vaghoo, H.; Surampudi, V.; Kultyshev, R.; Mandal, M.; Rasul, G.; Mathew, T.; Olah, G. A. *J. Org. Chem.* **2006**, *71*, 6806 and references cited therein.
9. Marchand, A. P. *Aldrichimica Acta*, **1995**, *28*, 95.
10. Krishnamurti, R.; Bellow, D. R.; Prakash, G. K. S. *J. Org. Chem.* **1991**, *56*, 984.
11. (a) Sasaki, T.; Eguchi, S.; Kiriyama, T.; Hiroaki, O. *Tetrahedron* **1974**, *30*, 2707. (b) Marchand, A. P.; Arney, B. E. Jr.; Dave, P. R.; Satyanarayana, N. *J. Org. Chem.*, **1988**, *53*, 1088. (c) Liebenberg, W.; Van der Walt, J. J.; Van der Schyf, C. J. *Pharmazie* **2000**, *55*, 833. (d) Marchand, A. P.; Keith, J. M.; Alihodzic, S.; Ganguly, B.; Somers, A.; W. Hariprakasha, H. K.; Power, T. D.; Watson W. H.; Bodige, S. G. *Struct. Chem.* **2001**, *12*, 313.
12. (a) Romański, J.; Młostów, G. *Synthesis*, **2002**, 1355. (b) Romański, J.; Marchand, A. P. *Pol. J. Chem.*, **2004**, *78*, 223.
13. Linden, A.; Romański, J.; Młostów, G.; Heimgartner, H. *Acta Crystallogr. Sect. C*, **2005**, *C61*, 221.
14. Quast, H.; Becker, Ch.; Witzel, M.; Peters, E.-M.; Peters K.; Schnering, H. G. von, *Liebigs Ann. Org. Bioorg. Chem* **1996**, *6*, 985.
15. Singh, R. P.; Twamley, B.; Shreeve, J. M. *J. Fluorine Chem.* **2001**, *112*, 329.
16. (a) Romański, J.; Młostów, G.; Żmudzka, K. *47<sup>th</sup> Annual Meeting of Polish Chemical Society*, Wrocław, Poland, September 12-17, 2004, Materials, K012, 58; (b) Romański, J. *Polish J. Chem.* **2007**, *81*, 187-191.
17. Prakash, G. K. S.; Yudin, A. K. *Chem. Rev.*, **1997**, *97*, 757.
18. Wenkert E.; Yoder, J. E. *J. Org. Chem.*, **1970**, *35*, 2986.
19. Bott, S. G.; Rajagopal, D.; Marchand, A. P.; Kumar, K. A. *J. Chem. Cryst.*, **1996**, *26*, 425.
20. Sasaki, T.; Eguchi, S.; Kiriyama, T.; Hiroaki, O. *Tetrahedron*, **1974**, *30*, 2707.
21. (a) Martins, F. J. C.; Viljoen, A. M.; Kruger, H. G.; Fourie, L.; Roscher, J.; Joubert, A. J.; Wessels, P. L. *Tetrahedron*, **2001**, *57*, 1601. (b) Romański, J.; Młostów, G.; Linden, A.; Heimgartner, H. *Pol. J. Chem.*, **2005**, *79*, 973.