

A facile synthesis of 3-aryl-2-mercaptoproacrylic acid under microwave irradiation

Jian-Feng Zhou,^{a,b*} Feng-Xia Zhu,^b Hui-Qin Zhu,^b and Yu-Lan Zhu,^{a,b}

^aDepartment of Chemistry, Huaiyin Teachers College, Huai'an 223300, P. R. China

^bJiangsu Key Laboratory for Chemistry of Low-Dimensional Materials,
Huai'an 223300, P. R. China

E-mail: jfzhou@163.com

Abstract

A series of 3-aryl-2-mercaptoproacrylic acid derivatives were synthesized by the hydrolysis of 5-(benzylidene)rhodanines in alkaline solution under microwave irradiation. Compared with the classical reaction conditions, this method has the advantage of excellent yields (75~94%), short reaction time (3~6 min), environmental benign, low energy consumption and easy work-up.

Keywords: Rhodanine, benzylidenerhodanines, 3- aryl-2-mercaptoproacrylic acids, microwave irradiation


Introduction

The study of 3-aryl-2-mercaptoproacrylic acids as analytical reagents has shown that they act as bidentate ligands and form colored complexes with transition metal ions with great sensitivity and which are easily extractable with organic solvents.^{1, 2} The 3-aryl-2-mercaptoproacrylic acids can inhibit of neuraminidases.³ They were also used as intermediates for the synthesis of a variety of heterocyclic sulphur compounds.⁴ The synthesis of 3-aryl-2-mercaptoproacrylic acids derivatives were generally carried out by the hydrolysis of 5-(benzylidene)rhodanines in alkaline solution in refluxing with longer reactions time (0.5~1.0hs).^{5, 6} To the best of our knowledge, although the synthesis of 3-aryl-2-mercaptoproacrylic acids derivatives via the hydrolysis of 5-(benzylidene)-rhodanines in alkaline solution has been studied previously^{1, 3, 5, 6}, such a kind of reaction has not been investigated under microwave irradiation until now.

The wide applicability of microwave irradiation in chemical reaction enhancement is due to high reaction rates, formation of cleaner products, and operational simplicity.^{7, 8, 9, 10} Continuing our interest in the development of efficient reagent and simple procedures for the synthesis of organic compounds, we have reported previously a facile synthesis of 5-(arylalkylidene) rhodanines in the presence of tetrabutylammonium bromide as a phase transfer catalyst in water

under microwave irradiation.¹¹ Herein, we would like to report an efficient synthesis of 3-aryl-2-mercaptopropanoic acid derivatives which were synthesized by the hydrolysis of 5-(benzylidene)rhodanines in alkaline solution under microwave irradiation (Scheme 1).

The synthetic route is shown as following



Ar: (a) C₆H₅, (b) 4-CH₃OC₆H₄, (c) 2-furyl, (d) C₆H₅CH=CH, (e) 4-NO₂C₆H₄, (f) 4-CH₃C₆H₄, (g) 2-ClC₆H₄, (h) 4-ClC₆H₄, (i) 4-(CH₃)₂NC₆H₄, (j) 3, 4-(OCH₂O)C₆H₃.

Scheme 1

Results and Discussion

As is shown in Scheme 1, the hydrolysis of 5-(benzylidene)rhodanines in alkaline solution under microwave irradiation to give 3-aryl-2-mercaptopropanoic acid **4a** in 91% yield. The results (summarized in Table 1) clearly reveal the scope and generality of the reaction with respect to various substituted aromatic aldehyde, and that reasonable yields (75~94%) of the products were achieved after only 3~6 min of microwave irradiation. Another important feature of this procedure is the survival of a variety of functional groups, such as ethers, nitro groups, halides, amino, heterocycles *etc.* under the reaction conditions. All products were characterized by IR, ¹H NMR, MS spectral data and Elemental analysis. Their melting points (except **4e~4j**) are identical to those of the known compounds reported in the literature.

Table 1. Synthesis of **4a~j** under microwave irradiation

Compds	Ar	Time / min	Yields / %*	Mp (°)	Lit. mp.(°)
3a	C ₆ H ₅	5	91	135~136	(133~134) ⁵
3b	4-CH ₃ OC ₆ H ₄	6	77	140~142	140 (dec) ⁶
3c	2-furyl	3	89	115~116	(115~117) ⁵
3d	C ₆ H ₅ CH=CH	5	82	150~152	148-151 ⁵
3e	4-NO ₂ C ₆ H ₄	3	94	140~142	--
3f	4-CH ₃ C ₆ H ₄	5	78	179~180	--
3g	2-ClC ₆ H ₄	3	80	140-143	--
3h	4-ClC ₆ H ₄	3	85	161-162	--
3i	4-(CH ₃) ₂ NC ₆ H ₄	6	75	163~164	--
3j	3,4-(OCH ₂ O)C ₆ H ₃	5	81	168~170	--

*Yields of the isolated products.

In conclusion, an efficient, simple procedure for the synthesis of 3-aryl-2-mercaptoproacrylic acid derivatives from the corresponding the hydrolysis of 5-(benzylidene)rhodanines in alkaline solution under microwave irradiation has been developed.

Experimental Section

General Procedures. All compounds prepared were characterized by ¹H NMR, IR and element analyses and were described in the experimental section. Melting points were determined in XT-5 digital melting point instrument and uncorrected. IR spectra were recorded on a Nicolet Avatar 360 FT-IR spectrometer. ¹H NMR spectra were measured at a Burke 400 MHz NMR spectrometer using TMS as internal standard and DMSO-d₆ as solvent. MS spectra were obtained on a LCQ Advanbage instrument. Elemental analyses were determined using Perkin-Elmer 240C Elemental Analysis. The reactions were carried out with a modified commercial microwave oven (Sanle WP650D 650w) under atmospheric pressure. 5-(benzylidene)rhodanines were prepared according to the literature.¹¹ Other chemicals were available commercially.

General experimental procedure. A mixture of the 5-(benzylidene)rhodanines (2mmol), 15 % sodium hydroxide (10 mL) in an Erlenmeyer flask (25ml) equipped with reflux condenser were irradiated in a microwave oven for 3~6 minutes (as indicated by TLC). The mixture was then cooled to room temperature. The concentrated hydrochloride acid was added. The precipitate formed was filtered out and recrystallized from ethanol.

- 4a.** M.p. 135~136°; ^1H NMR (400 MHz, DMSO-d6) δ 7.95(s, 1H, CH=), 7.43(s, 5H, Ar-H), 4.67 (s, 1H, SH); IR (KBr, cm^{-1}): 3043, 2566, 1665, 1593, 1505, 1447, 1180, 801; LC/MS (ESI): m/e = 179(M-H) $^-$; Anal. Calcd. for $\text{C}_9\text{H}_8\text{O}_2\text{S}$: C 59.98; H 4.47; Found: C 60.12; H 4.35.
- 4b.** M.p. 140~142°; ^1H NMR (400 MHz, DMSO-d6) δ 7.96 (s, 1H, CH=), 7.69(d, 2H, J = 8.8Hz, Ar-H), 6.98 (d, 2H, J = 8.8Hz, Ar-H), 4.65(s, 1H, SH), 3.84 (s, 3H, CH_3O); IR (KBr, cm^{-1}): 3103, 2999, 2568, 1655, 1588, 1508, 1439, 1176, 805; LC/MS (ESI): m/e = 209 (M-H) $^-$; Anal. Calcd. for $\text{C}_{10}\text{H}_{10}\text{O}_3\text{S}$: C 57.13; H 4.78; Found: C 57.19; H 4.56.
- 4c.** M.p. 115~116°; ^1H NMR (400 MHz, DMSO-d6) δ 8.23 (s, 1H, CH=), 7.80~7.42 (m, 3H, furyl-H), 4.69 (s, 1H, SH); IR (KBr, cm^{-1}): 3124, 2560, 1669, 1580, 1505, 1423, 1182, 802; LC/MS (ESI): m/e = 169 (M-H) $^-$; Anal. Calcd. for $\text{C}_7\text{H}_6\text{O}_3\text{S}$: 49.40; H 3.55; Found: C 49.52; H 3.46.
- 4d.** M.p. 150~152°; ^1H NMR (400 MHz, DMSO-d6) δ 7.97(d, 1H, 8.2 Hz, CH=), 6.95~7.44(m, 7H, Ar-H, -CH=CH-), 4.89 (s, 1H, SH); IR (KBr, cm^{-1}): 3126, 2601, 1675, 1610, 1560, 1425, 1180, 805; LC/MS (ESI): m/e = 205 (M-H) $^-$; Anal. Calcd. for $\text{C}_{11}\text{H}_{10}\text{O}_2\text{S}$: C 64.07; H 4.89; Found: C 64.21 H 4.76.
- 4e.** M.p. 140~142°; ^1H NMR (400 MHz, DMSO-d6) δ 7.91 (s, 1H, CH=), 8.31(d, 2H, J = 8.8Hz, Ar-H), 7.82 (d, 2H, J = 8.8Hz, Ar-H), 4.90(s, 1H, SH); IR (KBr, cm^{-1}): 3113, 2560, 1678, 1592, 1517, 1424, 1178, 800; LC/MS (ESI): m/e = 224 (M-H) $^-$; Anal. Calcd. for $\text{C}_9\text{H}_7\text{NO}_4\text{S}$: C 47.80; H 3.13; Found: C 47.90; H 3.05.
- 4f.** M.p. 179~180°; ^1H NMR (400 MHz, DMSO-d6) δ 7.91 (s, 1H, CH=), 7.78(d, 2H, J = 8.0Hz, Ar-H), 7.12 (d, 2H, J = 8.0Hz, Ar-H), 4.65 (s, 1H, SH), 2.36(s, 3H, CH_3); IR (KBr, cm^{-1}): 3088, 2919, 2566, 1670, 1592, 1505, 1418, 1185, 801; LC/MS (ESI): m/e = 193(M-H) $^-$; Anal. Calcd. for $\text{C}_{10}\text{H}_{10}\text{O}_2\text{S}$: C 61.83; H 5.19; Found: C 61.98; H 5.07
- 4g.** M.p. 140~143°; ^1H NMR (400 MHz, DMSO-d6) δ 7.87 (s, 1H, CH=), 7.75~7.43(m, 4H, Ar-H), 4.73 (s, 1H, SH); IR(KBr, cm^{-1}): 3058, 2561, 1678, 1589, 1508, 1430, 1182, 802; LC/MS(ESI): m/e = 213 (M-H) $^-$; Anal. Calcd. for $\text{C}_9\text{H}_7\text{ClO}_2\text{S}$: C 55.95; H 3.29; Found: C56.09; H 3.21.
- 4h.** M.p. 161~162°; ^1H NMR (400 MHz, DMSO-d6) δ 7.82 (s, 1H, CH=), 7.62 (d, 2H, J = 8.4Hz, Ar-H), 7.43 (d, 2H, J = 8.4Hz, Ar-H), 4.73 (s, 1H, SH); IR (KBr, cm^{-1}): 3103, 2572, 1664, 1586, 1505, 1422, 1180, 802°LC/MS(ESI): m/e = 213 (M-H) $^-$; Anal. Calcd. for $\text{C}_9\text{H}_7\text{ClO}_2\text{S}$: C 55.95; H 3.29; Found: C 56.08; H 3.22.
- 4i.** M.p. 163~164° ^1H NMR (400 MHz, DMSO-d6) δ 7.99 (s, 1H, CH=), 7.81(d, 2H, J = 8.8Hz, Ar-H), 7.52(d, 2H, J = 8.8Hz, Ar-H), 4.73 (s, 1H, SH), 3.15 (s, 6H, $\text{N}(\text{CH}_3)_2$); IR (KBr, cm^{-1}): 3124, 2575, 1695, 1513, 1436, 1182, 805; LC/MS(ESI): m/e = 222 (M-H) $^-$; Anal. Calcd. for $\text{C}_{11}\text{H}_{13}\text{NO}_2\text{S}$: C 59.17; H 5.89; Found: C 59.29; H 5.74.
- 4j.** M.p. 168~170°; ^1H NMR (400 MHz, DMSO-d6) δ 7.81 (s, 1H, CH=), 7. 52-7.27 (m, 3H, Ar-H), 4.77 (s, 1H, SH), 5.99 (s, 2H, OCH_2O); IR (KBr, cm^{-1}): 3083, 2905, 2821, 2576, 1654, 1579, 1500, 1449, 1180, 801; LC/MS (ESI): m/e = 223 (M-H) $^-$; Anal. Calcd. for $\text{C}_{10}\text{H}_8\text{O}_4\text{S}$: C 53.56; H 3.60;. Found: C 53.68; H 3.52.

Acknowledgements

We thank the National Natural Science Foundation of China (No. 20571029) and Jiangsu Key Laboratory for Chemistry of Low-Dimensional Materials for financial support.

References

1. Izquierdo, A.; Carrasco, J. *Analyst* **1984**, *109*, 605.
2. Izquierdo, A.; Beltrán, J. L. *Mikrochimica Acta* **1989**, *98*(1-3), 91.
3. Castineiras, A.; Gil, M. J.; Sevillano, P. *J. Molecular Structure* **2000**, *522*, 193.
4. Harhash, A. H.; Elnagdi, M. H.; Abdallah, S. O. *Indian J. Chem.* **1973**, *11*, 128.
5. Campaigne, E.; Cline, R. E. *J. Org. Chem.* **1956**, *21*, 32.
6. Shi H.; Wang, Z.; Shi, H. *Huaxue Shiji* **1995**, *17*, 329; *Chem. Abstr.*, **1996**, *124*, 316658.
7. Caddick, S. *Tetrahedron* **1995**, *51*, 10403.
8. Galema, S. A. *Chem. Soc. Rev.* **1997**, *26*, 233.
9. El Ashry, El Sayed H.; Ahmed, A. K. *Arkivoc* **2006**, (ix), 1.
10. Kappe, C. O. *Angew. Chem. Inter. Ed. Engl.* **2004**, *43*, 6250.
11. Zhou, J-F.; Zhu, F-X.; Song Y-Z.; Zhu, Y-L. *Arkivoc* **2006**, (xiv), 175.