

Supplementary Material

Pseudo-enantiomeric coupling reagents for predictable incorporation into the peptide chain D and/or L amino acid residue of racemic substrates

Justyna Fraczyk, Marta Michalska, Beata Kolesinska and Zbigniew J. Kaminski*

Institute of Organic Chemistry, Lodz University of Technology, Poland

Email: zbigniew.kaminski@p.lodz.pl

Table of Contents

Syntheses of dipeptides 14b-g using 2DMT/quinine/2BF₄⁻ (4) as coupling reagent	S2
S1. Synthesis of Z-D-Ala-Leu-OMe (14b).....	S2
S2. Synthesis of Z-D-Ala-Tyr-OMe (14c).....	S2
S3. Synthesis of Z-D-Ala-Ser-OMe (14d).....	S2
S4. Synthesis of Z-D-Ala-Val-OMe (14e).....	S2
S5. Synthesis of Z-D-Ala-Ala-OMe (14f).....	S2
S6. Synthesis of Z-D-Ala-Gly-OMe (14g).....	S2
Syntheses of dipeptides 15b-g using 2DMT/quinine/2BF₄⁻ (4) as coupling reagent	S2
S7. Synthesis of Z-Phe-Leu-OMe (15b).....	S2
S8. Synthesis of Z-Phe-Tyr-OMe (15c).....	S2
S9. Synthesis of Z-Phe-Ser-OMe (15d).....	S2
S10. Synthesis of Z-Phe-Val-OMe (15e).....	S3
S11. Synthesis of Z-Phe-Ala-OMe (15f).....	S3
S12. Synthesis of Z-Phe-Gly-OMe (15g).....	S3
Syntheses of dipeptides 14a-g using 2DMT/quinidine/2BF₄⁻ (7) as coupling reagent	S3
S13. Synthesis of Z-Ala-Phe-OMe (14a).....	S3
S14. Synthesis of Z-Ala-Leu-OMe (14b).....	S3
S15. Synthesis of Z-Ala-Tyr-OMe (14c).....	S3
S16. Synthesis of Z-Ala-Ser-OMe (14d).....	S3
S17. Synthesis of Z-Ala-Val-OMe (14e).....	S3
S18. Synthesis of Z-Ala-Ala-OMe (14f).....	S4
S19. Synthesis of Z-Ala-Gly-OMe (14g).....	S4
Syntheses of dipeptides 15b-f using 2DMT/quinidine/2BF₄⁻ (7) as coupling reagent	S4
S20. Synthesis of Z-D-Phe-Leu-OMe (15b).....	S4
S21. Synthesis of Z-D-Phe-Tyr-OMe (15c).....	S4
S22. Synthesis of Z-D-Phe-Ser-OMe (15d).....	S4
S23. Synthesis of Z-D-Phe-Val-OMe (15e).....	S4
S24. Synthesis of Z-D-Phe-Ala-OMe (15f).....	S4
Supporting References	S5

Syntheses of 14b-g using 2DMT/quinine/2BF₄⁻ (4) as coupling reagent.**S1.** Synthesis of Z-D-Ala-Leu-OMe (14b) using 2DMT/quinine/2BF₄⁻ (4) as coupling reagent.

In the synthesis according to typical procedure 2 were used: 2DMT/quinine/2BF₄⁻ (0.780 g, 1 mmol), quinine (0.101 g, 0,31 mmol), *rac*-Z-Ala-OH (0.446 g, 2 mmol), NMM (0.110 mL, 1 mmol) and HCl*H-Leu-OMe (0.182 g, 1 mmol). Z-D-Ala-Leu-OMe (0.214 g, yield 61%) was obtained, mp 68-70 °C, lit. mp 72-73 °C.^{S1}

GC: R_t 8.33 (L-Leu); R_t 3.54 (L-Ala); R_t 3.41 (D-Ala); L-Ala/D-Ala= 1/99.

S2. Synthesis of Z-D-Ala-Tyr-OMe (14c) using 2DMT/quinine/2BF₄⁻ (4) as coupling reagent.

In the synthesis according to typical procedure 2 were used: 2DMT/quinine/2BF₄⁻ (0.780 g, 1 mmol), quinine (0.101 g, 0,31 mmol), *rac*-Z-Ala-OH (0.446 g, 2 mmol), NMM (0.110 mL, 1 mmol) and HCl*H-Tyr-OMe (0.232 g, 1 mmol). Z-D-Ala-Tyr-OMe (0.228 g, yield 57%) was obtained, mp 102-105 °C, lit. mp 105-107 °C.^{S2}

GC: R_t 23.12 (L-Tyr); R_t 4.65 (L-Ala); R_t 3.53 (D-Ala); L-Ala/D-Ala= 0.5/99.5.

S3. Synthesis of Z-D-Ala-Ser-OMe (14d) using 2DMT/quinine/2BF₄⁻ (4) as coupling reagent.

In the synthesis according to typical procedure 2 were used: 2DMT/quinine/2BF₄⁻ (0.780 g, 1 mmol), quinine (0.101 g, 0,31 mmol), *rac*-Z-Ala-OH (0.446 g, 2 mmol), NMM (0.110 mL, 1 mmol) and HCl*H-Ser-OMe (0.156 g, 1 mmol). Z-D-Ala-Ser-OMe (0.191 g, yield 59%) was obtained, mp 128-130 °C, lit. mp 128-130 °C.^{S3}

GC: R_t 7.29 (L-Ser); R_t 3.56 (L-Ala); R_t 3.35 (D-Ala); L-Ala/D-Ala= 2/98.

S4. Synthesis of Z-D-Ala-Val-OMe (14e) using 2DMT/quinine/2BF₄⁻ (4) as coupling reagent.

In the synthesis according typical procedure 2 were used 2DMT/quinine/2BF₄⁻ (4) (0.780 g, 1 mmol), quinine (0.101 g, 0.31 mmol), *rac*-Z-Ala-OH (0.446 g, 2 mmol), NMM (0.110 mL, 1 mmol), HCl*Val-OMe (0.168 g, 1 mmol). Z-D-Ala-Val-OMe (0.202 g, yield 60%) was obtained, mp 85-87 °C, lit mp 79-82 °C.^{S3}

GC: R_t 4.76 (L-Val); R_t 3.56 (L-Ala); R_t 3.35 (D-Ala); L/D = 0.5/99.5.

S5. Synthesis of Z-D-Ala-Ala-OMe (14f) using 2DMT/quinine/2BF₄⁻ (4) as coupling reagent .

In the synthesis according to typical procedure 2 were used: 2DMT/quinine/2BF₄⁻ (0.780 g, 1 mmol), quinine (0.101 g, 0,31 mmol), *rac*-Z-Ala-OH (0.446 g, 2 mmol), NMM (0.110 mL, 1 mmol) and HCl*H-Ala-OMe (0.140, 1 mmol). Z-D-Ala-Ala-OMe (0.191, yield 62%) was obtained, mp 85-87 °C, lit. mp 105-106 °C.^{S4}

GC: R_t 3.57 (L-Ala); R_t 3.31 (D-Ala); L/D = 50.5/49.5.

S6. Synthesis of Z-D-Ala-Gly-OMe (14g) using 2DMT/quinine/2BF₄⁻ (4) as coupling reagent .

In the synthesis according to typical procedure 2 were used: 2DMT/quinine/2BF₄⁻ (0.780 g, 1 mmol), quinine (0.101 g, 0,31 mmol), *rac*-Z-Ala-OH (0.446 g, 2 mmol), NMM (0.110 mL, 1 mmol) and HCl*H-Gly-OMe (0.125 g, 1 mmol). Z-D-Ala-Gly-OMe (0.165 g, yield 56%) was obtained, mp 90-91 °C, lit. mp 94-96 °C.^{S5}

GC: R_t 4.82 (Gly); R_t 3.69 (L-Ala); R_t 3.41 (D-Ala); L-Ala/D-Ala= 2/98.

Syntheses of 15b-g using 2DMT/quinine/2BF₄⁻ (4) as coupling reagent.**S7.** Synthesis of Z-Phe-Leu-OMe (15b) using 2DMT/quinine/2BF₄⁻ (4) as coupling reagent.

In the synthesis according to typical procedure 2 were used: 2DMT/quinine/2BF₄⁻ (0.780 g, 1 mmol), quinine (0.101 g, 0,31 mmol), *rac*-Z-Phe-OH (0.599 g, 2 mmol), NMM (0.110 mL, 1 mmol) and HCl*H-Leu-OMe (0.182 g, 1 mmol). Z-Phe-Leu-OMe (0.260 g, yield 61%) was obtained, mp 109-112 °C, lit. mp 111 °C.^{S6}

GC: R_t 8.36 (L-Leu); R_t 18.72 (L-Phe); R_t 18.19 (D-Phe); L-Phe/D-Phe = 99.5/0.5.

S8. Synthesis of Z-Phe-Tyr-OMe (15c) using 2DMT/quinine/2BF₄⁻ (4) as coupling reagent.

In the synthesis according to typical procedure 2 were used: 2DMT/quinine/2BF₄⁻ (0.780 g, 1 mmol), quinine (0.101 g, 0,31 mmol), *rac*-Z-Phe-OH (0.599 g, 2 mmol), NMM (0.110 mL, 1 mmol) and HCl*H-Tyr-OMe (0.232 g, 1 mmol). Z-Phe-Tyr-OMe (0.272 g, yield 57%) was obtained, mp 154-157 °C, lit. mp 158 °C.^{S7}

GC: R_t 23.10 (L-Tyr); R_t 18.71 (L-Phe); R_t 18.14 (D-Phe); L-Phe/D-Phe = 99/1.

S9. Synthesis of Z-Phe-Ser-OMe (15d) using 2DMT/quinine/2BF₄⁻ (4) as coupling reagent.

In the synthesis according to typical procedure 2 were used: 2DMT/quinine/ 2BF_4^- (0.780 g, 1 mmol), quinine (0.101 g, 0,31 mmol), *rac*-Z-Phe-OH (0.599 g, 2 mmol), NMM (0.110 mL, 1 mmol) and $\text{HCl}^*\text{H-Ser-OMe}$ (0.156 g, 1 mmol). Z-Phe-Ser-OMe (0.236 g, yield 59%) was obtained, mp 86-77 °C, lit. mp 83-84 °C.^{s8}

GC: R_t 7.20 (L-Ser); R_t 18.51 (L-Phe); R_t 18.14 (D-Phe); L-Phe/D-Phe = 98/2.

S10. Synthesis of Z-Phe-Val-OMe (**15e**) using 2DMT/quinine/ 2BF_4^- (**4**) as coupling reagent.

In the synthesis according to typical procedure 2 were used: 2DMT/quinine/ 2BF_4^- (0.780 g, 1 mmol), quinine (0.101 g, 0,31 mmol), *rac*-Z-Phe-OH (0.599 g, 2 mmol), NMM (0.110 mL, 1 mmol) and $\text{HCl}^*\text{H-Val-OMe}$ (0.168 g, 1 mmol). Z-Phe-Val-OMe (0.247 g, yield 60%) was obtained, mp 110-113 °C, lit. mp 114-115 °C.^{s9}

GC: R_t 4.76 (L-Val); R_t 18.58 (L-Phe); R_t 18.14 (D-Phe); L-Phe/D-Phe = 98/2.

S11. Synthesis of Z-Phe-Ala-OMe (**15f**) using 2DMT/quinine/ 2BF_4^- (**4**) as coupling reagent.

In the synthesis according to typical procedure 2 were used: 2DMT/quinine/ 2BF_4^- (0.780 g, 1 mmol), quinine (0.101 g, 0,31 mmol), *rac*-Z-Phe-OH (0.599 g, 2 mmol), NMM (0.110 mL, 1 mmol) and $\text{HCl}^*\text{H-Phe-OMe}$ (0.139 g, 1 mmol). Z-Phe-Ala-OMe (0.238 g, yield 62%) was obtained, mp 121-123 °C, lit. mp 122-125 °C.^{s10}

GC: R_t 3.64 (L-Ala); R_t 18.58 (L-Phe); R_t 18.15 (D-Phe); L-Phe/D-Phe = 99/1.

S12. Synthesis of Z-Phe-Gly-OMe (**15g**) using 2DMT/quinine/ 2BF_4^- (**4**) as coupling reagent.

In the synthesis according to typical procedure 2 were used: 2DMT/quinine/ 2BF_4^- (0.780 g, 1 mmol), quinine (0.101 g, 0,31 mmol), *rac*-Z-Phe-OH (0.599 g, 2 mmol), NMM (0.110 mL, 1 mmol) and $\text{HCl}^*\text{H-Gly-OMe}$ (0.125 g, 1 mmol). Z-Phe-Gly-OMe (0.207 g, yield 56%) was obtained, mp 120-121 °C, lit. mp 120 °C.^{s11}

GC: R_t 4.74 (Gly); R_t 18.58 (L-Phe); R_t 18.14 (D-Phe); L-Phe/D-Phe = 97/3.

Syntheses of 14b-g using 2DMT/quinidine/ 2BF_4^- (7**) as coupling reagent.**

S13. Synthesis of Z-Ala-Phe-OMe (**14a**) using 2DMT/quinidine/ 2BF_4^- (**7**) as coupling reagent.

In the synthesis according to typical procedure 3 were used: 2DMT/quinidine/ 2BF_4^- (0.780 g, 1 mmol), quinidine (0.101 g, 0,31 mmol), *rac*-Z-Ala-OH (0.446 g, 2 mmol), NMM (0.110 mL, 1 mmol) and $\text{HCl}^*\text{H-Phe-OMe}$ (0.216 g, 1 mmol). Z-Ala-Phe-OMe (0.210 g, yield 55%) was obtained, mp 100-102 °C. lit. mp 103-104 °C.^{s12}

GC: R_t 18.71 (L-Phe); R_t 3.54 (L-Ala); R_t 3.30 (D-Ala); L-Ala/D-Ala = 99.5/0.5.

S14. Synthesis of Z-Ala-Leu-OMe (**14b**) using 2DMT/quinidine/ 2BF_4^- (**7**) as coupling reagent.

In the synthesis according to typical procedure 3 were used: 2DMT/quinidine/ 2BF_4^- (0.780 g, 1 mmol), quinidine (0.101 g, 0,31 mmol), *rac*-Z-Ala-OH (0.446 g, 2 mmol), NMM (0.110 mL, 1 mmol) and $\text{HCl}^*\text{H-Leu-OMe}$ (0.182 g, 1 mmol). Z-Ala-Leu-OMe (0.210 g, yield 60%) was obtained, mp 69-71 °C. lit. mp 72-73 °C.^{s13}

GC: R_t 18.71 (L-Phe); R_t 3.54 (L-Ala); R_t 3.30 (D-Ala); L-Ala/D-Ala = 99.5/0.5.

S15. Synthesis of Z-Ala-Tyr-OMe (**14c**) using 2DMT/quinidine/ 2BF_4^- (**7**) as coupling reagent.

In the synthesis according to typical procedure 3 were used: 2DMT/quinidine/ 2BF_4^- (0.780 g, 1 mmol), quinidine (0.101 g, 0,31 mmol), *rac*-Z-Ala-OH (0.446 g, 2 mmol), NMM (0.110 mL, 1 mmol) and $\text{HCl}^*\text{H-Leu-OMe}$ (0.182 g, 1 mmol). Z-Ala-Tyr-OMe (0.240 g, yield 60%) was obtained, mp 132-134 °C. lit. mp 122 °C.^{s14}

GC: R_t 23.28 (L-Tyr); R_t 4.61 (L-Ala); R_t 3.58 (D-Ala); L-Ala/D-Ala = 99.5/0.5.

S16. Synthesis of Z-Ala-Ser-OMe (**14d**) using 2DMT/quinidine/ 2BF_4^- (**7**) as coupling reagent.

In the synthesis according to typical procedure 3 were used: 2DMT/quinidine/ 2BF_4^- (0.780 g, 1 mmol), quinidine (0.101 g, 0,31 mmol), *rac*-Z-Ala-OH (0.446 g, 2 mmol), NMM (0.110 mL, 1 mmol) and $\text{HCl}^*\text{H-Ser-OMe}$ (0.156 g, 1 mmol). Z-Ala-Ser-OMe (0.188 g, yield 58%) was obtained, mp 130-132 °C. lit. mp 134-135 °C.^{s15}

GC: R_t 7.27 (L-Ser); R_t 3.55 (L-Ala); R_t 3.35 (D-Ala); L-Ala/D-Ala = 99/1.

S17. Synthesis of Z-Ala-Val-OMe (**14e**) using 2DMT/quinidine/ 2BF_4^- (**7**) as coupling reagent.

In the synthesis according to typical procedure 3 were used: 2DMT/quinidine/ 2BF_4^- (0.780 g, 1 mmol), quinidine (0.101 g, 0,31 mmol), *rac*-Z-Ala-OH (0.446 g, 2 mmol), NMM (0.110 mL, 1 mmol) and $\text{HCl}^*\text{H-Val-OMe}$ (0.168 g, 1 mmol). Z-Ala-Val-OMe (0.209 g, yield 62%) was obtained, mp 81-83 °C. lit. mp 82-84 °C.^{s16}

GC: R_t 4.75 (L-Val); R_t 3.55 (L-Ala); R_t 3.35 (D-Ala); L-Ala/D-Ala = 99/1.

S18. Synthesis of Z-Ala-Ala-OMe (**14f**) using 2DMT/quinidine/ 2BF_4^- (**7**) as coupling reagent.

In the synthesis according to typical procedure 3 were used: 2DMT/quinidine/ 2BF_4^- (0.780 g, 1 mmol), quinidine (0.101 g, 0,31 mmol), *rac*-Z-Ala-OH (0.446 g, 2 mmol), NMM (0.110 mL, 1 mmol) and $\text{HCl}^*\text{H-Ala-OMe}$ (0.139 g, 1 mmol). Z-Ala-Ala-OMe (0.188 g, yield 61%) was obtained, mp 109-113 °C. lit. mp 109-112 °C.^{s17}

GC: R_t 3.55 (L-Ala); R_t 3.33 (D-Ala); L-Ala/D-Ala = 99.5/0.5.

S19. Synthesis of Z-Ala-Gly-OMe (**14g**) using 2DMT/quinidine/ 2BF_4^- (**7**) as coupling reagent.

In the synthesis according to typical procedure 3 were used: 2DMT/quinidine/ 2BF_4^- (0.780 g, 1 mmol), quinidine (0.101 g, 0,31 mmol), *rac*-Z-Ala-OH (0.446 g, 2 mmol), NMM (0.110 mL, 1 mmol) and $\text{HCl}^*\text{H-Gly-OMe}$ (0.125 g, 1 mmol). Z-Ala-Gly-OMe (0.168 g, yield 57%) was obtained, mp 95-97 °C. lit. mp 97-98 °C.^{s18}

GC: R_t 4.85 (Gly); R_t 3.67 (L-Ala); R_t 3.40 (D-Ala); L-Ala/D-Ala = 99.5/0.5.

Syntheses of 15b-g using 2DMT/quinidine/ 2BF_4^- (7**) as coupling reagent.**

S20. Synthesis of Z-D-Phe-Leu-OMe (**15b**) using 2DMT/quinidine/ 2BF_4^- (**7**) as coupling reagent.

In the synthesis according to typical procedure 3 were used: 2DMT/quinidine/ 2BF_4^- (0.780 g, 1 mmol), quinidine (0.101 g, 0,31 mmol), *rac*-Z-Phe-OH (0.599 g, 2 mmol), NMM (0.110 mL, 1 mmol) and $\text{HCl}^*\text{H-Leu-OMe}$ (0.182 g, 1 mmol). Z-D-Phe-Leu-OMe (0.268 g, yield 63%) was obtained, mp 139-141 °C. lit. mp 124-125 °C.^{s19}

GC: R_t 8.35 (L-Leu); R_t 18.70 (L-Phe); R_t 18.15 (D-Phe); L-Phe/D-Phe = 2/98.

S21. Synthesis of Z-D-Phe-Tyr-OMe (**15c**) using 2DMT/quinidine/ 2BF_4^- (**7**) as coupling reagent.

In the synthesis according to typical procedure 3 were used: 2DMT/quinidine/ 2BF_4^- (0.780 g, 1 mmol), quinidine (0.101 g, 0,31 mmol), *rac*-Z-Phe-OH (0.599 g, 2 mmol), NMM (0.110 mL, 1 mmol) and $\text{HCl}^*\text{H-Tyr-OMe}$ (0.232 g, 1 mmol). Z-D-Phe-Tyr-OMe (0.310 g, yield 65%) was obtained, mp 159-163 °C. lit. mp 131-132 °C for Z-Phe-Tyr-OMe.^{s20}

GC: R_t 23.12 (L-Tyr); R_t 18.70 (L-Phe); R_t 18.15 (D-Phe); L-Phe/D-Phe = 0.5/99.5.

S22. Synthesis of Z-D-Phe-Ser-OMe (**15d**) using 2DMT/quinidine/ 2BF_4^- (**7**) as coupling reagent.

In the synthesis according to typical procedure 3 were used: 2DMT/quinidine/ 2BF_4^- (0.780 g, 1 mmol), quinidine (0.101 g, 0,31 mmol), *rac*-Z-Phe-OH (0.599 g, 2 mmol), NMM (0.110 mL, 1 mmol) and $\text{HCl}^*\text{H-Ser-OMe}$ (0.182 g, 1 mmol). Z-D-Phe-Ser-OMe (0.156 g, yield 39%) was obtained, mp 145-147 °C. lit. mp 146-147 °C.^{s21}

GC: R_t 7.20 (L-Ser); R_t 18.50 (L-Phe); R_t 18.14 (D-Phe); L-Phe/D-Phe = 1/99.

S23. Synthesis of Z-D-Phe-Val-OMe (**15e**) using 2DMT/quinidine/ 2BF_4^- (**7**) as coupling reagent.

In the synthesis according to typical procedure 3 were used: 2DMT/quinidine/ 2BF_4^- (0.780 g, 1 mmol), quinidine (0.101 g, 0,31 mmol), *rac*-Z-Phe-OH (0.599 g, 2 mmol), NMM (0.110 mL, 1 mmol) and $\text{HCl}^*\text{H-Val-OMe}$ (0.168 g, 1 mmol). Z-D-Phe-Val-OMe (0.247 g, yield 60%) was obtained, mp 99-101 °C. 139-132 °C.^{s22}

GC: R_t 4.75 (L-Val); R_t 18.55 (L-Phe); R_t 18.15 (D-Phe); L-Phe/D-Phe = 1/99.

S24. Synthesis of Z-D-Phe-Ala-OMe (**15f**) using 2DMT/quinidine/ 2BF_4^- (**7**) as coupling reagent.

In the synthesis according to typical procedure 3 were used: 2DMT/quinidine/ 2BF_4^- (0.780 g, 1 mmol), quinidine (0.101 g, 0,31 mmol), *rac*-Z-Phe-OH (0.599 g, 2 mmol), NMM (0.110 mL, 1 mmol) and $\text{HCl}^*\text{H-Ala-OMe}$ (0.139 g, 1 mmol). Z-D-Phe-Ala-OMe (0.238 g, yield 62%) was obtained, mp 141-143 °C. lit. mp 135-135.5 °C.^{s23}

GC: R_t 3.64 (L-Ala); R_t 18.58 (L-Phe); R_t 18.15 (D-Phe); L-Phe/D-Phe = 0.5/99.5.

Supporting References

1. Polglase, W. J.; Smith, E. L., *J. Am. Chem. Soc.*, **1949**, *71*, 3083-3084.
2. Izumiya, N; Fruton, J. S. *J. Biol. Chem.* **1956**, *218*, 59-76.
3. Kolesinska, B.; Kaminski, Z. *J. Org. Letters*, **2009**, *11*, 765-768.
4. Kaminski, Z. J.; Kolesinska, B.; Kaminska, J. E.; Gora, J. *J. Org. Chem.* **2001**, *66*, 6276-6281.
5. Hammadi, A.; Nuzillard, J. M.; Poulin, J. C.; Kagan, H. B. *Tetrahedron Asymm.* **1992**, *3*, 1247-1262.
6. Singh, U.; Ghosh, S. K.; Chadha, M. S.; Mamdapur, V. R. *Tetrahedron Lett.*, **1991**, *32*, 255-258.
7. Sun, Q.; Xu, B.; Niu, Y.; Xu, F.; Liang, L.; Wang, C.; Yu, J.; Jin, H.; Xu, P. *ChemMedChem*, **2015**, *10*, 498-510.
8. Ranganathan, D.; Shah, K.; Vaish, N. *J. Chem. Soc. Chem. Commun.* **1992**, *16*, 1145-1147.
9. Kraznobrizhii, H. R.; Skrynuk, E. M.; Kowalenko, L. G. *J. Gen. Chem. USSR. (Engl. Transl.)* **1975**, *45*, 1381-1383.
10. Trzuppek, L. S.; Go, A.; Kopple, K. D. *J. Org. Chem.* **1979**, *44*, 4577-4581.
11. Romani, S.; Moroder, L. Bavermann, G.; Wuensch, E. *Synthesis* **1985**, *8*, 738-742.
12. Kawasaki, K.; Tsuji, T.; Maeda, M.; Matsumoto, T.; Hirase, K. *Chem. Pharm. Bull.* **1987**, *35*, 1044-1048.
13. Chen, H.; He, M.; Wang, Y.; Zhai, L.; Cui, Y.; Li, Y.; Lee, Y.; Zhou, H.; Hong, X.; Deng, Z. *Green Chem.* **2011**, *13*, 2723-2726.
14. Neuenschwander, M.; Fahrni, H. P.; Lienhard, U. *Helv. Chim. Acta*, **1978**, *61*, p. 2437-2451.
15. Marchiori, F.; Rocchi, R.; Moroder, L.; Fontana, A.; Scoffone, E., *J. Am. Chem. Soc.* **1968**, *90*, 5885-5889.
16. Palomo, C.; Aizpurua, J. M.; Loinaz, I.; Fernandez-Berridi, M. J.; Irusta, L. *Org. Lett.* **2001**, *3*, 2361-2364.
17. Iida, Y.; Sanekata, F. *Chem. Pharm. Bull.* **1990**, *38*, 2997-3003.
18. Matoba, H. *Agricult. Biol. Chem.* **1972**, *36*, 1423-1425.
19. Moersch, G. W. et al. *J. Med. Chem.* **1979**, *22*, 935-943.
20. Isowa, Y.; Ichikawa, T. *Bull. Chem. Soc. Japan*, **1979**, *52*, 796-800.
21. Nakajima, K.; Kawai, H.; Takai, M.; Okawa, K. *Bull. Chem. Soc. Jpn.* **1977**, *50*, 917-920.
22. Sokolowska, T.; Biernat, J. F. *Roczniki Chem.* **1966**, *40*, 1665-1673.
23. Carpino, L. A.; Chao, H. G.; Tien, J-H. *J. Org. Chem.* **1989**, *54*, 4302-4313.